

AD-A237 133



2

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1235

R & T Code 4133020

Technical Report No.14

Discreteness-of-Solvent Effects in Marcus Equations in the Mean Spherical Approximation

by

L. Blum and W.R. Fawcett

Prepared for Presentation

at

The Electrochemical Society Meeting

Washington D.C., May 1991

Department of Chemistry  
University of California  
Davis, CA 95616

May 20, 1991

Reproduction in whole or in part is permitted  
for any purpose of the United States Government

"This document has been approved for public release  
and sale; its distribution is unlimited"

DTIC  
COPY  
INSPECTED  
5

by  
Classification  
Availability Codes  
Dist Special

A-1

~~91 6 18 122~~

91 6 18 124

91-02543



REPORT DOCUMENTATION PAGE			Form Approved MAY 1962 EDITION
May 20, 1991		Technical	
Discreteness-of-Solvent Effects in Marcus Equations in the Mean Spherical Approximation		N00014-90-J-1235	
L. Blum and W.R. Fawcett			
Department of Chemistry University of California Davis, CA 95616		No. 14	
Office of Naval Research 800 N. Quincy Arlington, VA 22217-5000			
Prepared for Presentation at The Electrochemical Society Meeting, Washington D.C., May 1991.			
Unclassified			
<p>In the Marcus theory of electron transfer the outer sphere contribution to the free energy of activation <math>G_{os}</math> is usually computed from Born continuum dielectric formula. The Mean Spherical Approximation gives systematic corrections to this quantity, which for a solvent of hard dipolar spheres can be explicitly calculated. The contributions are of two kinds: The polarization of the solvent by the charges, and the electrostriction effects, which usually are small. the effects of ion association are included using recent results for sticky potential models. These corrections also can be incorporated into the dynamical theories of solvation effects.</p>			
Gibbs solvation energies of ions, outer sphere reorganization energy for electron transfer		5	
unclassified		unclassified	
unclassified		unclassified	

# 75-Word Abstract Form

Extended Abstract must be submitted with the 75-Word Abstract by December 1, 1990

## Washington, DC, May 5-10, 1991

Submit to: The Electrochemical Society, Inc.  
10 South Main Street, Pennington, NJ 08534-2836  
With a copy to the Organizer

Abstract No.  
(to be assigned by the Society)

Schedule for  
Title of Symposium: **Electrode Processes V**

Sponsored by  
Division/Group: **Physical Electrochemistry**

Title of paper: **The Discreetness of Solvent Effects in Marcus Equations in the Mean Spherical Approximation**

Authors: **L. BLUM, DEPARTMENT OF PHYSICS, UNIVERSITY OF PUERTO RICO, RIO PIEDRAS, PR 00931-3343, USA**

Business affiliation and complete mailing address: **and W.R. FAWCETT, DEPARTMENT OF CHEMISTRY, UNIVERSITY OF CALIFORNIA, DAVIS CA 95616**

State or Country:

(ZIP Code)

(Telephone No.)

(Type abstract in this area—double-spaced.)

**The Discreetness of Solvent Effects in Marcus Equations in the Mean Spherical Approximation:** *L. Blum*, Department of Physics, University of Puerto Rico, Rio Piedras, PR 00931-3343, USA and *W.R. Fawcett*, Department of Chemistry, University of California, Davis CA 95616.

In the Marcus theory of electron transfer the outer sphere contribution to the free energy of activation  $G_{os}$  is usually computed from Born continuum dielectric formula. The Mean Spherical Approximation gives systematic corrections to this quantity, which for a solvent of hard dipolar spheres can be explicitly calculated. The contributions are of two kinds: The polarization of the solvent by the charges, and the electrostriction effects, which usually are small. The effects of ion association are included using recent results for sticky potential models. These corrections also can be incorporated into the dynamical theories of solvation effects.

Do you require any audiovisual equipment?

☐ 35 mm (2 x 2 in.) slide projector

☒ Overhead projector

☐ Other equipment at author's expense and subject to availability

Has the information in this abstract been presented verbally, submitted for publication, or published?

☐ Yes ☒ No

If the answer is yes, please provide the reference (except in the case of invited review presentations).

Is a full length paper on this work to be submitted for Society Journal publication? ☐ Yes ☐ No

The Extended Abstracts will be published in a volume copyrighted by The Electrochemical Society, Inc.

The Discreteness of Solvent Effects in Marcus  
Equations in the Mean Spherical Approximation:

L. Blum, Department of Physics, University of Puerto  
Rico, Rio Piedras, PR 00931-3343, USA and W.R.

Fawcett, Department of Chemistry, University of  
California, Davis, CA 95616

In the Marcus theory [1] of electron transfer the outer  
sphere contribution to the free energy of activation  $\Delta G_{os}$  is  
usually computed from Born continuum dielectric formula.  
The Mean Spherical Approximation gives systematic  
corrections to this quantity for molecular solvents [2].  
For a solvent of hard dipolar spheres it can be explicitly  
calculated. The contributions are of two kinds [3]: The  
polarization of the solvent by the charges, and the  
electrostriction effects,

$$\Delta G_{os} = \Delta G_{os}^{is} + \Delta G_{os}^{ss} \quad (1)$$

$$\Delta G_{os}^{is} = - \frac{2z_1^2 e^2 (1 - 1/\epsilon)}{\sigma_1 + \sigma_s \lambda} \quad (2)$$

the second term is more complex, and can be found in the  
literature.  $\sigma_1$  is the ionic diameter and  $\sigma_s$  is the solvent  
diameter. The parameter  $\lambda$  is obtained from the dielectric  
constant  $\epsilon$  using [4]

$$\lambda(1 + \lambda)^2 = 4\sqrt{\epsilon} \quad (3)$$

and is related to reaction field at infinite coupling (dipole  
moment). The influence of ion association is included  
using a sticky potential approximation.

When the dipole diameter,  $\sigma_s$ , vanishes, then the classic  
Born approximation is obtained.

Calculations have been carried out for a representative  
group of solvents. These calculations are discussed with  
respect to data in the literature.

References

- [1] R. Marcus, *J. Chem. Phys.* 43, 679 (1965).
- [2] L. Blum, *J. Stat. Phys.* 18, 451 (1978).
- [3] F. Garisto, P. Kusalik and G.N. Patey, *J. Chem. Phys.* 79, 6294 (1983).
- [4] M.S. Wertheim, *J. Chem. Phys.* 55, 4291 (1971).